



# LASIK POSTOP

Hoopes Vision  
11820 S. State Street  
Draper, UT 84020  
801.568.0200



**ANTIBIOTIC:**  
Moxifloxacin/Ofloxacin  
**EVERY 4 HOURS  
FOR 1 WEEK**

Wait at least  
5 minutes  
between drops.  
Only necessary  
during waking  
hours.

**ARTIFICIAL TEARS  
(PRESERVATIVE FREE)**



**1-2 Drops EVERY 15-30 MIN  
THE DAY OF SURGERY  
THEN  
AT LEAST EVERY HOUR  
FOR 1 MONTH**



**STEROID:**  
Prednisolone Acetate  
**EVERY 2 HOURS  
UNTIL YOUR 1 DAY POSTOP VISIT  
THEN  
EVERY 4 HOURS  
FOR 1 WEEK**

\* SHAKE BEFORE  
EACH USE

## AFTER YOUR SURGERY:


- A short nap is advised. Limit activity and phone used the day of surgery. Increase activity as your comfort and vision allows. Take your normal over-the-counter pain medication if needed.
- Your vision will be foggy for the first day.
- You may experience some tearing, burning, stinging, light sensitivity, redness, and foreign body sensation.
- Do not drive until after your 1-day post-op appointment. You may return to work and/or drive when you are confident with your vision and your comfort allows.
- If you notice increasing pain or a sudden decrease in vision – call us immediately.

## ARTIFICIAL TEARS:

- FIRST MONTH – Use preservative free artificial tears in vials 1-2 drops **every hour**
- MONTH 2 & 3 - OK to switch to a bottled artificial tear and use every 2 to 3 hours.

## OTHER IMPORTANT PRECAUTIONS:

- Wash and dry your hands before putting in your eye drops. Be careful not to touch the tip of the bottle to any part of your eye. You may notice an unusual taste after using some drops.
- Your vision may fluctuate for the first several weeks.
- You may experience halos and glare around lights at night for the first couple weeks.
- No eye makeup for 3 - 5 days. New mascara is recommended after the procedure.
- Avoid getting soap or water in your eyes while showering for the first week.
- Wear your shields while sleeping or napping for the first week.
- You should protect your eyes from bright sun and wind for 1 month.
- No swimming or hot tubs for 2 weeks. No high-impact water sports for 3 months.
- Do not rub or bump your eyes for at least 2 months after surgery. It is a bad habit to rub your eyes. We suggest you avoid this habit altogether.



Preservative Free  
Artificial Tears (vials)  
Month 1



OK to use Bottle  
Artificial Tears  
when used less  
frequently

**Hoopes Vision Phone Number: 801-568-0200**

**Benjamin Buckner, MD**

Patient Signature: \_\_\_\_\_

Date: \_\_\_\_\_

Staff Signature: \_\_\_\_\_

Tomorrow's Appointment Time: